

The GEOMED 2 project: A high resolution geoid of the Mediterranean Sea

The GEOMED 2 research group

The GEOMED 2 partnership

Riccardo Barzaghi, Alberta Albertella, Daniela Carrion

Politecnico di Milano, Italy (Polimi)

Sean Bruinsma, Sylvain Bonvalot

GET UMR 5563, Toulouse, France (GET)

Marie-Françoise Lequentrec-Lalancette, Corinne Salaun

SHOM, Brest, France (SHOM)

Pascal Bonnefond

OCA/Géoazur, Sophia-Antipolis, France (OCA)

Georgios Vergos, Ilias Tziavos, Vassilios Grigoriadis

Aristotle University of Thessaloniki, Greece (AUTH)

Per Knudsen, Ole Andersen

DTU Space, Copenhagen, Denmark (DTU)

Mehmet Simav, Hasan Yildiz

General Command of Mapping, Ankara, Turkey (GCM)

Tomislav Basic, Matej Varga, Olga Bjelotomic

University of Zagreb, Zagreb, Croatia (UZ)

Antonio J. Gil

University of Jaén, Jaén, Spain (UJA)

The GEOMED 2 project (1/2)

- The project aims at estimating the geoid over the area


$$30 < \varphi < 48 \quad -10 < \lambda < 40$$

with a resolution of $3' \times 3'$

- ***Gravity data only will be used in the computation***
- Methods to be used in the geoid computation:
 - i) Collocation
 - ii) Stokes
 - iii) Stokes-FFT

The GEOMED 2 project (2/2)

- DOT will be then computed by difference with existing MSS altimetry derived estimates (e.g. the DTU2013)



The DTU2013 MSS over the Mediterranean Sea

- The circulation in the Mediterranean Sea will be also estimated

The Adriatic Sea test

- The selected area

$$36 < \varphi < 48 \quad 10 < \lambda < 21$$

- Gravity data selected with a mean spacing of $2' \times 2'$ from the following databases:

- i) BGI
- ii) SHOM
- iii) Croatia
- iv) Greece
- v) Italy
- vi) EGM2008 (void areas along the Eastern edge)

- Geoid estimate based on the Remove-Restore method

- The DTM/bathymetry model is **SRTM/NOAA** bathymetry at $3'' \times 3''$ (bathymetry is originally at $1' \times 1'$) in the window
$$35 < \varphi < 49 \quad 9 < \lambda < 22$$
- The **GGN GOCE-DIR5** to d/o **230** was used for the long wavelength modelling of gravity/geoid
- RTC effect was computed using the **GRAVSOFT TC** program ($r=80$ km from each computation point, data point heights equal to DTM). The reference DTM was estimated by low-pass filtering the detailed DTM (using a $25'$ cap)
- Residual gravity values were gridded using the **GRAVSOFT GEOGRID** program on a regular $3' \times 3'$ geographical grid in the area

$$36 < \varphi < 48 \quad 10 < \lambda < 21$$


The statistics of the remove step

	Δg_{obs}	$\Delta g_R = \Delta g_{\text{obs}} - \Delta g_{\text{MOD}}$	$\Delta g_r = \Delta g_{\text{obs}} - \Delta g_{\text{MOD}} - \Delta g_{\text{RTC}}$
n	82657	82657	82657
E(mgal)	11.398	-4.835	-0.853
σ (mgal)	46.676	32.205	20.786
min(mgal)	-184.800	-245.144	-166.528
MAX(mgal)	267.710	216.964	119.615

The statistics of the gridded 3'×3' residual gravity



	$\Delta g_r(\text{GRID})$
n	53261
E(mgal)	-0.304
σ (mgal)	17.799
min(mgal)	-136.760
MAX(mgal)	104.801

The residual gravity field



The applied geoid estimation methods

- i) FastCollocation (Bottini and Barzaghi, 1993)
- ii) Collocation (Lequentrec-Lalancette et al., 2002)
- iii) 1D_FFT (Haagmans et al., 1993)
- iv) 2D_FFT (Strang van Hess, 1990)



The residual geoid statistics

	N_r (FastCol)	N_r (Collocation)	N_r (1D_FFT)	N_r (2D_FFT)
n	53261	53261	53261	53261
E(m)	-0.148	-0.005	-0.200	-0.181
$\sigma(m)$	0.801	0.468	0.930	0.924
min(m)	-2.742	-2.267	-3.173	-3.136
MAX(m)	1.720	1.429	1.928	1.971

The statistics of the differences between N_r estimate: 1D_FFT vs FastCollocation

	$N_r(1D_FFT) - N_r(FastCollocation)$
n	22701
E(m)	-0.061
$\sigma(m)$	0.117


(In the area $38 < \varphi < 46$ $12 < \lambda < 19$)

The statistics of the differences between N_r estimates: 1D_FFT vs 2D_FFT

	$N_r(1D_FFT) - N_r(2D_FFT)$
n	22701
E(m)	-0.019
$\sigma(m)$	0.016

(In the area $38 < \varphi < 46$ $12 < \lambda < 19$)

The differences between N_r estimates: 1D_FFT vs FastCollocation



The statistics of the differences w.r.t. GPS/lev points

$$v = (N_{grav} - N_{GPS/lev}) - DN(q, l) = (N_{grav} - N_{GPS/lev}) - dx \sin q \cos l + dy \sin q \sin l + dz \cos q$$



$$q = 90^\circ - j ; \quad N_{GPS/lev} = h - H \quad (dx, dy, dz) \text{ datum shift translation}$$

Italy (mainland)


	v(FASTCOL)	v(1DFFT)	v(EGM2008)
n	594	590	595
E(m)	0.000	0.000	0.000
$\sigma(m)$	0.095	0.084	0.084
min(m)	-0.243	-0.294	-0.227
MAX(m)	0.260	0.239	0.276

Greece

	v(FASTCOL)	v(1DFFT)	v(EGM2008)
n	126	126	126
E(m)	0.000	0.000	0.000
$\sigma(m)$	0.181	0.164	0.118
min(m)	-0.489	-0.485	-0.304
MAX(m)	0.547	0.545	0.434



The DOT in the Adriatic Sea based on the DTU2013 MSS and the 1D_FFT geoid estimate



Comments and conclusions

- The Adriatic Sea Test proved that further checks on the gravity data are needed (consistency among different gravity databases)
- The processing chain seems to be satisfactory but some refinements must be implemented (possibly a denser gravity database should be selected based on a 1'x1' selection grid)
- The GOCE-DIR5 GGM allows an effective data reduction; other solutions will be also tested in the future (e.g. the time-wise and the space-wise solutions)
- Differences in the geoid computation methods are quite large and, to some extent, unexpected (ad hoc analysis will be devised to come to more coherent solutions)